

Research in Astrophysics from Space (E)  
Origin of Cosmic Rays (E1.6)  
Consider for oral presentation.

## **ANOMALOUS TRANSPORT OF HIGH ENERGY COSMIC RAYS IN GALACTIC SUPERBUBBLES**

Nasser Barghouty, [abdulnasser.f.barghouty@nasa.gov](mailto:abdulnasser.f.barghouty@nasa.gov)  
NASA Marshall Space Flight Center, Huntsville, Alabama, Alabama, United States

High-energy cosmic rays may exhibit anomalous transport as they traverse and are accelerated by a collection of supernovae explosions in a galactic superbubble. Signatures of this anomalous transport can show up in the particles' evolution and their spectra. In a continuous-time-random-walk (CTRW) model assuming standard diffusive shock acceleration theory (DSA) for each shock encounter, and where the superbubble (an OB stars association) is idealized as a heterogeneous region of particle sources and sinks, acceleration and transport in the superbubble can be shown to be sub-diffusive. While the sub-diffusive transport can be attributed to the stochastic nature of the acceleration time according to DSA theory, the spectral break appears to be an artifact of transport in a finite medium. These CTRW simulations point to a new and intriguing phenomenon associated with the statistical nature of collective acceleration of high energy cosmic rays in galactic superbubbles.